



Suitable and unsuitable micro-organisms

The following lists are selected micro-organisms which present minimum risk given good practice. As a result of changes to the hazard categorisation of certain micro-organisms by the Advisory Committee on Dangerous Pathogens (ACDP)¹, these tables supersede the lists found in the CLEAPSS *Laboratory Handbook* (1992), *Microbiology: An HMI Guide for Schools and Further Education* (1990), *Topics in Safety* (1988) and *Safety in Science Education* (1996).

Additionally, the tables include points of educational use and interest and comment on growing and maintaining the cultures. The lists of micro-organisms are not definitive; other organisms may be used if competent advice is obtained².

It should be noted that strains of micro-

organisms can differ physiologically and therefore may not give expected results. Where possible, fungi that produce large numbers of air-borne spores should be handled before sporulation occurs, so that the spread of spores into the air and possible risks of allergy or the triggering of asthmatic attacks are minimised. *This is particularly important for some genera, such as Aspergillus and Penicillium, which produce very large numbers of easily-dispersed spores.*

It should be noted that certain genera of these two fungi, previously listed as unsuitable for use in schools, are now not thought to present such a serious risk to health, given good practice in culture and handling.

References

1. Updates to the categories in this list are published from time to time and available only online as a standalone publication *The Approved List of Biological Agents*. See www.hse.gov.uk/pubns/misc208.pdf.

2. Organisations which can be consulted about the suitability of micro-organisms: Association for Science Education*; CLEAPSS*; Microbiology in Schools Advisory Committee; National Centre for Biotechnology Education; Scottish Schools Equipment Research Centre*.

(* Members only).

See 'Useful links' on www.misac.org.uk

Bacteria

Bacterium	Educational use/interest/suitability	Ease of use/maintenance
<i>Acetobacter aceti</i>	Of economic importance in causing spoilage in beers and wines. Oxidises ethanol to ethanoic (acetic) acid and ultimately to carbon dioxide and water.	Needs special medium and very frequent sub-culturing to maintain viability.
<i>Agrobacterium tumefaciens</i>	Causes crown galls in plants; used as a DNA vector in the genetic modification of organisms.	Grows on nutrient agar, but requires 2-3 days' incubation.
<i>Alcaligenes eutrophus</i>	In the absence of nitrogen, it produces intracellular granules of poly-β-hydroxybutyrate (PHB); was used in the production of biodegradable plastics.	Grows on nutrient agar.
<i>Azotobacter vinelandii</i>	A free-living nitrogen fixer, producing a fluorescent, water-soluble pigment when grown in iron (Fe)-limited conditions.	Grows on a nitrogen-free medium.
<i>Bacillus megaterium</i>	Has very large cells; produces lipase, protease and also PHB (see <i>Alcaligenes</i>); Gram-positive staining.	Grows on nutrient agar.

Bacterium	Educational use/interest/suitability	Ease of use/maintenance
<i>Bacillus stearothermophilus</i>	Thermophilic species which grows at 65°C; produces lipase and protease. Also used to test the efficiency of autoclaves.	Grows on nutrient agar.
<i>Bacillus subtilis</i> †	General-purpose, Gram-positive bacterium. Produces amylase, lipase and protease.	Grows on nutrient agar.
<i>Cellulomonas</i> sp.	Produces extra-cellular cellulase.	Grows on nutrient agar but also used with agar containing carboxymethylcellulose.
<i>Chromatium</i> species	A photosynthetic, anaerobic bacterium.	Requires special medium and light for good growth.
<i>Erwinia carotovora</i> (= <i>E. atroseptica</i>)	Produces pectinase which causes rotting in fruit and vegetables. Useful for studies of Koch's postulates.	Grows on nutrient agar.
<i>Escherichia coli</i> †	K12 strain: general-purpose, Gram-negative bacterium. B strain: susceptible to T4 bacteriophage.	Grows on nutrient agar.
<i>Janthinobacterium</i> (= <i>Chromobacterium lividum</i>)*	Produces violet colonies. Grows best at 20°C.	Needs frequent subculture and is best grown on glucose nutrient agar and broth.
<i>Lactobacillus</i> species	Ferment glucose and lactose, producing lactic acid; <i>L. bulgaricus</i> is used in the production of yoghurt.	Require special medium containing glucose and yeast extract and frequent subculturing to maintain viability.
<i>Leuconostoc mesenteroides</i>	Converts sucrose to dextran: used as a blood plasma substitute.	Requires special medium as for <i>Lactobacillus</i> .
<i>Methylophilus methylotrophus</i>	Requires methanol as energy source; was used for the production of 'Pruteen' single-cell protein.	Requires special medium containing methanol.
<i>Micrococcus luteus</i> (= <i>Sarcina lutea</i>)	Produces yellow colonies; useful in the isolation of the bacterium from impure cultures. Also used to simulate the effects of disinfectants, mouthwashes and toothpastes on more harmful organisms. General-purpose, Gram-positive bacterium.	Grows on nutrient agar.
<i>Photobacterium phosphoreum</i>	Actively-growing, aerated cultures show bioluminescence; grows in saline conditions.	Requires a medium containing sodium chloride.
<i>Pseudomonas fluorescens</i>	Produces a fluorescent pigment in the medium.	Grows on nutrient agar.
<i>Rhizobium leguminosarum</i>	A symbiotic, nitrogen fixer; stimulates the formation of nodules on the roots of legumes. Only fixes nitrogen in plants.	Grows on yeast malt agar; some authorities recommend buffering with chalk to maintain viability.
<i>Rhodospseudomonas palustris</i>	A photosynthetic, anaerobic, red bacterium. Also grows aerobically in the dark.	Requires light and a special medium, growing atypically on nutrient agar.
<i>Spirillum serpens</i>	Of morphological interest.	May grow on nutrient agar but requires very frequent subculturing to maintain viability.
<i>Staphylococcus albus</i> (<i>epidermidis</i>)**	A general-purpose, Gram-positive bacterium, producing white colonies.	Grows on nutrient agar.
<i>Streptococcus</i> (= <i>Enterococcus</i>) <i>faecalis</i>	Of morphological interest, forming pairs or chains of cocci.	Nutrient agar with added glucose can be used but grows better on special medium, as for <i>Lactobacillus</i> .
<i>Streptococcus</i> (= <i>Lactococcus</i>) <i>lactis</i>	Of morphological interest, forming pairs or chains of cocci. Commonly involved in the souring of milk; also used as a starter culture for dairy products.	Can grow on nutrient agar with added glucose; some authorities recommend buffering with chalk to maintain viability.

† Some strains have been associated with health hazards. Reputable suppliers should ensure that safe strains are provided.

* Can be chosen for investigations that once required the use of *Chromobacterium violaceum* or *Serratia marcescens*.

** This organism has been known to infect debilitated individuals and those taking immuno-suppressive drugs. Some authorities advise against its use.

Bacterium	Educational use/interest/suitability	Ease of use/maintenance
<i>Streptococcus thermophilus</i>	Ferments glucose and lactose, producing lactic acid; used in the production of yoghurt. Grows at 50°C.	Can grow on nutrient agar with added glucose; some authorities recommend frequent subculturing to maintain viability.
<i>Streptomyces griseus</i>	Responsible for the earthy odour of soil. Grows to form a fungus-like, branching mycelium with aerial hyphae bearing conidia. Produces streptomycin.	Grows on nutrient or glucose nutrient agar but better on special medium which enhances formation of conidia.
<i>Thiobacillus ferrooxidans</i>	Involved in the bacterial leaching of sulphur-containing coal. Oxidises iron(II) and sulphur. Demonstrates bacterial leaching of coal samples containing pyritic sulphur.	Requires special medium.
<i>Vibrio natriegens</i> * (= <i>Beneckea natriegens</i>)	A halophile, giving very rapid growth. Prone, however, to thermal shock with a sudden drop in temperature.	Requires medium containing sodium chloride.

Fungi

Fungus	Educational use/interest/suitability	Ease of use/maintenance
<i>Agaricus bisporus</i>	Edible mushroom; useful for a variety of investigations on factors affecting growth.	Grows on compost containing well-rotten horse manure; available as growing 'kits'.
<i>Armillaria mellea</i>	The honey fungus; causes decay of timber and tree stumps. Produces rhizomorphs.	Grows very well on malt agar. Some authorities recommend carrot agar.
<i>Aspergillus nidulans</i> *	For studies of nutritional mutants. Produces abundant, easily-dispersed spores - may become a major laboratory contaminant!	Grows on Czapek Dox yeast agar. Special media required for studying nutritional mutants.
<i>Aspergillus niger</i> *	Useful for studies of the influence of magnesium on growth and the development of spore colour. Used commercially for the production of citric acid. Produces abundant, easily-dispersed spores - may become a major laboratory contaminant!	Requires special sporulation medium for investigations.
<i>Aspergillus oryzae</i> *	Produces a potent amylase; useful for studies of starch digestion. Also produces protease. Used by the Japanese in the production of rice wine (saki).	Grows on malt agar; add starch (or protein) for investigations.
<i>Botrytis cinerea</i>	Causes rotting in fruits, particularly strawberries. Useful for studies of Koch's postulates with fruit, vegetables and <i>Pelargonium</i> sp. Important in the production of some dessert wines ('noble' rot). Used in ELISA protocols.	Can be grown on malt agar or agar with oatmeal.
<i>Botrytis fabae</i>	Causes disease in bean plants.	Requires agar with oatmeal.
<i>Candida utilis</i>	Simulates behaviour of pathogenic <i>Candida</i> spp. in investigations of fungicidal compounds.	Grows on malt agar or glucose nutrient agar.
<i>Chaetomium globosum</i>	Useful for studies of cellulase production; thrives on paper.	Can be grown on V8 medium but survives well just on double thickness wall paper, coated with a flour paste.
<i>Coprinus lagopus</i>	For studies of fungal genetics.	Grows on horse dung.
<i>Eurotium</i> (= <i>Aspergillus</i>) <i>repens</i>	Produces yellow cleistocarps (cleistothecia) embedded in the medium and green conidial heads in the same culture.	Requires special medium.
<i>Fusarium graminearum</i>	Causes red rust on wheat; used in the manufacture of 'Quorn' mycoprotein.	Can be grown on V8 medium.

* A well-known supplier currently lists an unspecified species of *Vibrio* because of its morphological interest. This has a typical shape, better shown than by *V. natriegens*. However, this species is a Hazard Group 2 organism which may cause human disease. This bacterium should only be used in establishments that have containment facilities suitable for work with Hazard Group 2 microorganisms.

* Possible risk of allergy/asthma if large numbers of spores are inhaled.

Fungus	Educational use/interest/suitability	Ease of use/maintenance
<i>Fusarium oxysporum</i>	A pathogen of many plants. Produces sickle-cell shaped spores, a red pigment and pectinase.	Grows well on several media including malt, potato dextrose and Czapek Dox yeast agar.
<i>Fusarium solani</i>	Digests cellulose; macroconidia have a sickle shape.	Grows on potato dextrose agar.
<i>Helminthosporium avenae</i>	A pathogen of oats.	May not grow easily in laboratory cultures.
<i>Kluyveromyces lactis</i>	A yeast, isolated from cheese and dairy products. Ferments lactose and used to convert dairy products to lactose-free forms. Genetically-modified strains are used to produce chymosin (rennet).	Grows on malt agar or glucose nutrient agar.
<i>Leptosphaeria maculans</i>	For studies of disease in Brassica plants.	Requires cornmeal agar or prune yeast lactose agar to promote sporulation in older cultures.
<i>Monilinia</i> (= <i>Sclerotinia</i>) <i>fructigena</i>	For studies of brown rot in apples. Useful for studies of Koch's postulates.	Grows on malt agar or potato dextrose agar.
<i>Mucor genevensis</i>	For studies of sexual reproduction in a homothallic strain of fungus.	Grows on malt agar.
<i>Mucor hiemalis</i>	For studies of sexual reproduction between heterothallic + and - strains and zygospore production.	Grows on malt agar.
<i>Mucor mucedo</i>	Common black 'pin mould' on bread. For sporangia (asexual), mating types and amylase production.	Grows on malt agar.
<i>Myrothecium verucaria</i>	For studies of cellulose decomposition but <i>Chaetomium globosum</i> is preferred.	Grows on malt agar.
<i>Neurospora crassa</i> *	Red bread mould. Produces different coloured ascospores. Can be used in studies of genetics. Beware - readily becomes a major laboratory contaminant!	Grows on malt agar.
<i>Penicillium chrysogenum</i> *	Produces penicillin; useful for comparative growth inhibition studies in liquid media or when inoculated on to agar plates seeded with Gram-positive and negative bacteria. Produces yellow pigment.	Grows on malt agar, though some authorities indicate that it thrives better on liquid media.
<i>Penicillium expansum</i> *	Does not produce penicillin; causes disease in apples. Useful for studies of Koch's postulates.	Grows on malt agar.
<i>Penicillium notatum</i> *	Produces penicillin; useful for comparative growth inhibition studies in liquid media or when inoculated onto agar plates seeded with Gram-positive and negative bacteria.	Grows on malt agar.
<i>Penicillium roqueforti</i> *	Does not produce penicillin; the familiar mould of blue-veined cheese.	Grows on malt agar.
<i>Penicillium wortmanii</i> *	Produce wortmin rather than penicillin.	Grows on malt agar.
<i>Phaffia rhodozyma</i>	A fermenting red yeast. Used to colour the food supplied to fish-farmed salmon.	Grows on yeast malt agar.
<i>Phycomyces blakesleanus</i>	Produces very long sporangiophores which are strongly phototropic.	Grows on malt agar.
<i>Physalospora obtusa</i>	An ascomycete fungus that grows on apple. Thought to produce pectinase.	Grows on potato dextrose agar.
<i>Phytophthora infestans</i> +	Causes potato blight. Produces motile zoospores.	Can be grown on V8 medium.

* Possible risk of allergy/asthma if large numbers of spores are inhaled.

+ Now classed as a protoctist, so may not be listed under fungi by some suppliers.

Fungi	Educational use/interest/suitability	Ease of use/maintenance
<i>Plasmodiophora brassicae</i>	For studies of disease in Brassica plants, particularly club root. Useful for studies of Koch's postulates.	May not grow easily in culture.
<i>Pleurotus ostreatus</i>	Edible oyster cap mushroom.	Can be grown on rolls of toilet paper!
<i>Pythium de baryanum</i> +	Causes 'damping off' of seedlings; cress is best to use.	Grows on cornmeal agar.
<i>Rhizopus oligosporus</i>	Used in the fermentation of soya beans to make 'tempe', a meat-substitute food in Indonesia.	Grows on potato dextrose agar, Czapek Dox yeast agar and other fungal media.
<i>Rhizopus sexualis</i>	Produces rhizoids and zygospores. Useful for studies of the linear growth of fungi.	Grows on potato dextrose agar and other fungal media.
<i>Rhizopus stolonifer</i>	Produces rhizoids. Produces lipase.	Grows on potato dextrose agar, potato carrot agar, Czapek Dox yeast agar and other fungal media.
<i>Rhytisma acerinum</i>	An indicator of air pollution: less common in industrial areas. On sycamore leaves, it forms 'tar' spot lesions, the number or diameter of which can be compared at different sites.	Difficult to maintain but laboratory cultures are not likely to be needed.
<i>Saccharomyces cerevisiae</i>	Valuable for work in baking and brewing, showing budding, for spontaneous mutation and mutation-induction experiments, and for gene complementation using adenine- and histidine-requiring strains.	Grows on malt agar or glucose nutrient agar.
<i>Saccharomyces diastaticus</i>	Able to grow on starch by producing glucoamylase.	Grows on malt agar and nutrient agar + 1% starch.
<i>Saccharomyces ellipsoideus</i>	Used in fermentations to produce wine; can tolerate relatively high concentrations of ethanol.	Grows on malt agar.
<i>Saprolegnia litoralis</i> +	Parasitic on animals. Produces zoospores. Good illustration of asexual and sexual stages.	Culture by baiting pond water with hemp seeds.
<i>Schizosaccharomyces pombe</i>	Large cells, dividing by binary fission. Good for studies of population growth, using a haemocytometer for cell counts. Prone to thermal shock.	Grows on malt agar. For studies of population growth, a malt extract broth can be used.
<i>Sordaria brevicollis</i>	For studies of fungal genetics, including inheritance of spore colour and crossing over in meiosis.	Requires special medium for crosses between strains.
<i>Sordaria fimicola</i>	For studies of fungal genetics, including inheritance of spore colour and crossing over in meiosis.	Grows on cornmeal, malt and other agars but may not transfer readily from one medium to another. White-spore strain may not always grow normally on standard cornmeal agar.
<i>Sporobolomyces sp.</i>	Found on leaf surfaces. Spores are ejected forcibly into the air from mother cells.	Grows on malt, yeast malt and glucose nutrient agar but laboratory cultures may not be needed.
<i>Trichoderma reesei</i>	Commercial production of cellulase.	Grows on malt agar.

+ Now classed as a protocist, so may not be listed under fungi by some suppliers.

Viruses

These are rarely used in schools and colleges but a selected list of those which might be considered is given below.

Bacteriophage (T type) (host E. coli)
Potato Virus X
Tobacco Mosaic Virus

Cucumber Mosaic Virus
Potato Virus Y (not the virulent strain)
Turnip Mosaic Virus

Algae, protozoa (including slime moulds) and lichens

Though some protozoa are known to be pathogenic, the species quoted for experimental work in recent science projects and those obtained from schools' suppliers or derived from hay infusions, together with species of algae and lichens, are acceptable for use in schools.

Unsuitable micro-organisms

A number of micro-organisms have in the past been suggested for use in schools but are no longer considered suitable; these are listed below. Some fungi previously considered unsuitable have been reinstated in the list of selected organisms now that it is thought that they do not present a major risk, given good practice.

Bacteria

Chromobacterium violaceum
Clostridium perfringens (welchii)
Pseudomonas aeruginosa
Pseudomonas solanacearum
Pseudomonas tabaci
Serratia marcescens
Staphylococcus aureus
Xanthomonas phaseoli

Fungi

Rhizomucor (Mucor) pusillus